



[www.info4PI.org](http://www.info4PI.org)

## 10 Warning Signs of **Primary Immune Deficiency** for Adults

Primary Immune Deficiency (PI) causes children and adults to have infections that come back frequently or are unusually hard to cure. 1:500 persons are affected by one of the known Primary Immune Deficiencies. **If you or someone you know is affected by two or more of the following Warning Signs, speak to a physician about the possible presence of an underlying Primary Immune Deficiency.**

***Baxter***

**1** Two or more new ear infections within 1 year.

**2** Two or more serious sinus infections within 1 year, in the absence of an allergy.

**3** One pneumonia per year for more than 1 year.

**4** Chronic diarrhea with weight loss.

**5** Recurrent viral infections (colds, herpes, warts, condyloma).

**6** Recurrent need for intravenous antibiotics to clear infections.

**7** Recurrent, deep abscesses of the skin or internal organs.

**8** Persistent thrush or fungal infection on skin or elsewhere.

**9** Infection with normally harmless tuberculosis-like bacteria.

**10** A family history of PI.

---

[www.ImmuneDisease.com](http://www.ImmuneDisease.com)

These warning signs were developed by the Jeffrey Modell Foundation Medical Advisory Board. Consultation with Primary Immune Deficiency experts is strongly suggested.

For information or referrals, contact the Jeffrey Modell Foundation at 866-INFO-4-PI or [info4PI.org](http://info4PI.org).

Baxter is a registered trademark of Baxter International Inc.

© 2010 Jeffrey Modell Foundation  
February 2010 HYL4379A 093311

***Baxter***